

Variations in avoidable hospital admissions and emergency visits by primary care characteristics: an ecological study in Italy

Zeno Dalla Valle¹, Giovanni Emanuele Ricciardi¹, Carlo Signorelli¹, Cristina Renzi^{1,2}

Keywords: Primary healthcare; healthcare outcomes; inappropriate hospitalisation; emergency presentations

Parole chiave: Assistenza sanitaria primaria; esiti sanitari; ospedalizzazione inadeguata

Abstract

Background. Primary healthcare plays a central role in providing preventive care, managing chronic conditions and reducing inappropriate emergency presentations. The study aimed at providing population-level evidence on the correlation between the characteristics of primary healthcare across Italian regions and health outcomes included in the National Programs Outcomes of the National Agency for Regional Healthcare Services.

Study Design. Ecological study

Methods. We analysed healthcare data from the National Agency for Regional Healthcare Services, the public lists of primary care doctors and the National Federation of Surgeons and Dentists and the National Institutes of Statistics referring to the 20 Italian regions. Pearson's correlation and Spearman's correlation were used to assess the relationships between primary healthcare characteristics and health outcomes.

Results. Overall, across all Italian regions each general practitioner had on average 1447 patients and was 57.5 years old. The study found positive correlations between the number of patients per general practitioner and non-urgent Emergency Department visits among adult patients (Pearson's $r = 0.58$, $p = 0.008$), the number of residents aged 65+ per general practitioner and the rate of chronic obstructive pulmonary disease admissions (Pearson's $r = 0.49$, $p = 0.029$), and the age of general practitioners and lower-extremity amputations in diabetes patients (Pearson's $r = 0.56$, $p = 0.011$). A negative correlation was observed between the age of general practitioners and urinary tract infection admissions (Pearson's $r = -0.76$; $p < 0.001$). A non-linear negative correlation was found between the age of general practitioners and chronic obstructive pulmonary disease admissions (Spearman's $\rho = -0.46$, $p = 0.041$).

Conclusions. The findings emphasise the importance of guaranteeing sufficient numbers of primary healthcare physicians to meet patients' needs, and for limiting avoidable hospitalisations and emergency presentations. General practitioners' age might also influence the provision of care, but more research is needed on possible mechanisms.

¹ Faculty of Medicine, University Vita-Salute San Raffaele, Milan, Italy

² Epidemiology of Cancer Healthcare & Outcomes (ECHO) Research Group, Department of Behavioural Science and Health, Institute of Epidemiology & Health Care, University College London, London UK

Introduction

Primary health care (PHC) is a whole-of-government and whole-of-society approach to health that combines three core components: multisectoral policy and action, empowered people and communities, and primary care and essential public health functions as the core of integrated health services (1,2). In Italy, PHC is provided by the "primary care doctors", divided into *general practitioners* (medici di medicina generale, GPs) for the adults and *primary care paediatricians* (pediatri di libera scelta, PLSs) for children. Primary care doctors are the first point of contact for patients, playing a crucial role in providing appropriate healthcare assistance, preventative healthcare, reducing the number of inappropriate specialist visits, Emergency Department (ED) and hospital outpatient attendance, etc. (3). Moreover, GPs are key healthcare providers for people with mental illness, with collaborative programmes having been developed between PHC and mental healthcare specialists, for example the "G. Leggieri Program" in Emilia-Romagna (4).

All citizens are registered with a primary care doctor, and sometimes between the age of 6 and 14 years, they must switch from a PLS to a GP, with only some exceptions (5). The current legislation and national collective contract of GPs and PLSs established 1,500 and 880 as the maximum number of patients each, while the optimal number is set at 1,000 and 600, respectively (5–7). Some regional and local health authorities raised those limits to address the shortage of primary care doctors and guarantee primary care to the entire population (8,9).

There is limited research on the relationship between primary care organization and health outcomes in Italy. The National Agency for Regional Healthcare Services (Agenzia Nazionale per i Servizi Sanitari Regionali, Agenas) annually releases the national report "Programma Nazionale Esiti" (National Outcomes Program, PNE), analyzing various healthcare indicators. One section focuses on primary care indicators, including non-urgent daytime emergency department visits and potentially avoidable hospital admissions (10,11). The aim of the present ecological study was to analyze the correlation between the characteristics of primary care across Italian regions and PNE indicators.

Methods

We obtained the list of GPs and PLSs working in Italy from various regional and national health

authorities' websites. We determined each doctor's age using data from the National Federation of Surgeons and Dentists (FNOMCeO) website. Due to varying website update schedules, we reported the data as available on each site (Supplementary Table 1). We used geographical and demographic data updated to 2022 from the National Institutes of Statistics (Istituto Nazionale di Statistica, ISTAT).

For each of the 20 Italian regions, we analysed the average age and the average number of residents assigned to each primary care doctor, the latter serving as an indirect measure of the number of registered patients per GP and PLS. This measure was calculated as the total number of residents in the region aged 14 years or older, and the total number of residents aged less than 14 years, respectively, divided by the number of GPs and PLSs in the region. For GPs, we also examined the number of patients aged 65 years or older and 75 years or older.

Finally, we investigated the correlation between such characteristics and several indicators from the PNE 2023 report (which is based on data from the year 2022). For GPs the indicators included are non-urgent ED visit rate of adult patients (≥ 14 years of age) during daytime (from 8 A.M. to 8 P.M.) from Monday to Friday (12), lower-extremity amputation among patients with diabetes rate (13), asthma among adults admission rate (14), chronic obstructive pulmonary disease (COPD) admission rate (15), diabetes short-term and long-term complication admission rate (16), uncontrolled diabetes (without complications) admission rate (17), urinary tract infection admission rate (18), influenza admission rate (19), hypertension admission rate (20), heart failure admission rate (21) and involuntary psychiatric treatment (22). For PLSs, the indicators included are non-urgent daytime ED visit rate among children (23), asthma among children admission rate (24) and paediatric acute gastroenteritis admission rate (25). We used values adjusted for age and sex, calculated by Agenas as described in every indicator protocol, and we aggregated them considering the average value for each region.

The dataset was formed by 20 observations, one for each region, with the characteristics of the primary healthcare doctors and the PNE indicators analysed.

The correlation between primary care characteristics and PNE outcome indicators was evaluated with the Pearson correlation coefficient and the Spearman correlation coefficient. For linear correlations, linear regression models were fitted too.

P-values < 0.05 were considered statistically significant. Analyses were conducted using Python v. 3.10.9

and the following libraries: pandas v. 1.5.3, scipy v. 1.10.1, statsmodels v. 0.13.5, matplotlib v. 3.7.1, seaborn v. 0.12.2. The library italy-geopop v. 0.6.2 was used to retrieve geographical and demographic data, as it includes data from 2022 from ISTAT.

Results

Overall, across all Italian regions each GP had on average 1,447 patients (range: 1,075-1,879) and 400 patients over-65 years (range across regions: 302-497). GPs were 57.5 years old on average (range across regions: 51.6-67.1). Each PLS had 1,139 patients on average (range: 837-1,905) and was 58.9 years old (range across regions: 53.6-66.0). PNE indicators description is reported in Tab. 1.

The study found a positive correlation between the number of patients per GP and the rate of non-urgent adult visits to emergency departments during the day (Pearson's $r = 0.58$, $p = 0.008$). Additionally, there was a positive correlation between the number of over-65 residents per GP and the rate of COPD admissions to hospitals (Pearson's $r = 0.49$, $p = 0.029$). Further, the study found a positive correlation between the age of GPs and the rate of lower-extremity amputations among patients with diabetes (Pearson's $r = 0.56$, $p = 0.011$). Conversely, a negative correlation was observed between the age of GPs and the rate of urinary tract infection admissions (Pearson's $r = -0.76$;

$p < 0.001$). Among with those findings, we fitted the linear regression models (Figure 1).

The study also reported a non-linear negative correlation between the age of GPs and COPD admission rate (Spearman's $\rho = -0.46$, $p = 0.041$). No significant correlations were found between the number of patients per PLS and any of the PNE indicators, nor between the age of PLSs and these indicators.

Discussion

Our study highlighted a positive correlation between the number of residents per GP and non-urgent ED visits, as well as between the number of over-65 residents per GP and COPD admission rate. Moreover, we found that age of GPs was positively correlated with lower-extremity amputation rates among patients with diabetes, while it was negatively correlated with urinary tract infection admissions.

A high number of patients per GP might lead to greater difficulties for patients in accessing primary healthcare with patients resorting for ED access. Previous Italian studies are scant, but a few studies reported a lower likelihood of ED visits for patients who have greater access to primary care medicine (26) and demonstrated that increasing primary care accessibility limited inappropriate ED use, estimating the magnitude of reduction to be 10% to 15% in inappropriate visits (27). This is also in line with available

Table 1. PNE indicators description.

	Mean (SD)	Min-Max
Non-urgent day time ED adults visit rate (%)	55.2 (17.3)	31.0 - 94.9
Lower-extremity amputation among patients with diabetes rate (x100,000)	7.6 (2.3)	4.8 - 12.8
Asthma among adults admission rate (x100,000)	3.2 (2.0)	1.0 - 8.0
COPD admission rate (x10,000)	13.4 (3.9)	7.7 - 20.8
Diabetes short-term and long-term complication admission rate (x100,000)	28.9 (8.7)	12.0 - 43.6
Uncontrolled diabetes (without complications) admission rate (x100,000)	6.8 (4.1)	0.0 - 17.0
Urinary tract infection admission rate (x100,000)	61.3 (39.3)	18.4 - 161.6
Influenza admission rate (x100,000)	6.2 (9.5)	0.5 - 41.5
Hypertension admission rate (100,000)	12.3 (5.3)	4.4 - 24.3
Heart failure admission rate (x10,000)	26.5 (4.8)	17.4 - 32.4
Involuntary psychiatric treatment rate (x10,000)	1.0 (0.5)	0.1 - 2.0
Non-urgent day time ED children visit rate (%)	77.2 (23.3)	42.7 - 126.6
Asthma among children admission rate (x10,000)	1.6 (0.9)	0.1 - 3.8
Paediatric acute gastroenteritis admission rate (x10,000)	4.9 (2.4)	1.4 - 10.6

PNE: programma nazionale esiti, ED: emergency department, COPD: chronic obstructive pulmonary disease, SD: standard deviation.




Figure 1. Correlations between the characteristics of GPs and the PNE indicators with the equations representing the fitted models. ED: emergency department, GP: general practitioner, COPD: chronic obstructive pulmonary disease.

studies from countries with a Beveridge-type National Healthcare System. In England, 26.5% of ED visits probably were preceded by the impossibility for patients to obtain a visit with the GP within a reasonable time (28). General practices providing more timely access has been shown to be associated with fewer ED visits per registered patient, however the number of GPs/1,000 patients was not independently associated with this outcome (29). In Sweden and Belgium the

most common pathway for emergency presentation is self-referred walk-in and only a small percentage of ED visits were a referral from a general practitioner (30,31). The main reason for non-urgent use of the ED might be that patients feel that it takes too long to wait for an appointment with their primary care doctor, even if they were successful in scheduling one (32). In contrast, other studies suggested that potentially avoidable ED visits were higher in deprived

areas, while they were not correlated with access to primary care (33).

COPD is a major cause of hospital admissions worldwide, especially in elderly patients who are the group with the highest number of prevalent cases (34). Effective primary care, including patient education, regular monitoring and therapeutic interventions can reduce COPD-related hospital admissions (35), but increasing number of older and complex patients per doctor might limit the possibilities of providing high quality care. Some previous studies have also suggested that rates of COPD admissions can be influenced by the prevalence of diagnosed COPD and socioeconomic status rather than other practice characteristics, such as practice performance scores (36).

The correlations between the age of GPs and PNE's outcomes are difficult to explain, as evidence on the topic is scant and further research would be needed to better understand the underlying mechanisms. While our analyses did not reveal a correlation between GP number and involuntary psychiatric treatment rate, more research on this is necessary, examining also additional process and outcome indicators for mental healthcare. Furthermore, it would be interesting to analyse the relationship between primary care characteristics and other healthcare indicators, including timely cancer diagnosis and cancer outcomes (37).

The findings of the present study should be considered in the context of several limitations. First, due to the ecological design we cannot establish causality, nor can we evaluate confounding factors. The sources we used to obtain the list of GPs and PLSs were not all up to date. Since we used Pearson's coefficient and Spearman's coefficient to assess the correlation, we might have missed other types of correlation (e.g. quadratic, etc.). Data from ISTAT let us calculate the number of residents per GP and PLS, <14 and ≥ 14 years respectively, based on the assumption that every resident, having the right to be assisted by a primary care doctor, is indeed registered with such a doctor. The number of residents per doctor is assumed to be a valid proxy indicator for the number of patients registered with a primary care doctor, but, as an aggregated average value, does not provide information about the variability. Moreover, primary care doctors can manually limit their number of patients before reaching the maximum number of patients. Agenas calculated non-urgent ED visits rate using white and green triage discharge colours as cases (12,23). The discharge triage colour is attributed evaluating the conditions at the moment of discharge, after the diagnostic and therapeutic process have been completed,

while the admission triage code could differ, reflecting more precisely whether the ED visit was urgent or not. More studies and a punctual analysis conducted at GP-level could be useful to further explore the findings and shed light on underlying mechanisms.

Conclusions

This ecological study explored the correlation between the characteristics of primary care doctors in Italy and various healthcare outcomes. The findings suggest that a higher patient load per GP is associated with an increased rate of non-urgent ED visits. Additionally, a higher proportion of over-65 residents per GP is linked to a higher rate of COPD admissions. Moreover, the age of GPs exhibits a complex relationship with healthcare outcomes. The study findings underscore the importance of optimizing primary care delivery to improve patient outcomes. Increasing the number of GPs and improving accessibility to primary care services can contribute to reducing inappropriate ED utilization and associated costs. More research is needed to better understand the causal mechanisms underlying these associations and to develop targeted interventions.

Riassunto

Variazione dei ricoveri e degli accessi in pronto soccorso potenzialmente evitabili in funzione delle caratteristiche delle cure primarie in Italia: uno studio ecologico

Introduzione. L'assistenza primaria è un componente cruciale dei sistemi sanitari, svolgendo un ruolo centrale nel fornire assistenza preventiva, gestire le condizioni croniche e ridurre le visite non urgenti al pronto soccorso. Questo studio ha lo scopo di indagare la correlazione tra le caratteristiche dei medici di medicina generale in Italia e gli indicatori sanitari riportati nel Piano Nazionale Esiti dell'Agenzia Nazionale per i Servizi Sanitari Regionali.

Disegno dello studio. Studio ecologico.

Metodi. abbiamo analizzato i dati del Programma Nazionale Esiti, le liste pubbliche di medici di medicina generale e pediatri di libera scelta e i dati della Federazione Nazionale degli Ordini dei Medici Chirurghi e Odontoiatri riferiti alle 20 regioni italiane. Abbiamo utilizzato i coefficienti di correlazione di Pearson e di Spearman per valutare le correlazioni.

Risultati. Complessivamente la media del numero di pazienti per medico di medicina generale era 1447 e l'età media dei medici era 57,5 anni. Lo studio ha trovato correlazioni positive tra il numero di pazienti per medico di base e il tasso di visite non urgenti al pronto soccorso tra gli adulti (r di Pearson = 0,58, $p = 0,008$), il numero di cittadini di età superiore ai 65 anni per medico di base e il tasso di

ricoveri per broncopneumopatia cronica ostruttiva (r di Pearson = 0,49, p = 0,029) e l'età dei medici di base e il tasso di amputazioni agli arti inferiori tra i pazienti con diabete (r di Pearson = 0,56, p = 0,011). Al contrario, è stata osservata una correlazione inversa tra l'età dei medici di base e il tasso di ricoveri per infezioni del tratto urinario (r di Pearson = -0,76; p < 0,001). È stata trovata una correlazione non lineare inversa tra l'età dei medici di base e il tasso di ricoveri per BPCO (ρ di Spearman = -0,46, p = 0,041).

Conclusioni. Questi risultati sottolineano l'importanza di garantire un numero sufficiente di medici per le cure primarie per soddisfare i bisogni di salute e limitare ospedalizzazioni potenzialmente evitabili e accessi potenzialmente impropri al pronto soccorso. Anche l'età dei medici di medicina generale potrebbe influenzare la fornitura di assistenza, ma sono necessarie ulteriori ricerche sui possibili meccanismi.

References

1. Declaration of Alma-Ata [Internet]. Available from: <https://iris.who.int/bitstream/handle/10665/347879/WHO-EURO-1978-3938-43697-61471-eng.pdf> [Last accessed: 2023 Dec 1].
2. A vision for primary health care in the 21st century: towards universal health coverage and the Sustainable Development Goals [Internet]. Geneva: World Health Organization and the United Nations Children's Fund (UNICEF); 2018 (Technical series on primary health care). Available from: <https://iris.who.int/bitstream/handle/10665/328065/WHO-HIS-SDS-2018.15-eng.pdf> [Last accessed: 2023 Dec 1].
3. Signorelli C. Igiene e sanità pubblica. Secrets. Domande e risposte. 3rd ed. SEU; 2021.
4. Berardi D, Ferrannini L, Menchetti M, Vaggi M. Primary care psychiatry in Italy. *J Nerv Ment Dis*. 2014 Jun;202(6):460–3.
5. ACCORDO COLLETTIVO NAZIONALE PER LA DISCIPLINA DEI RAPPORTI CON I MEDICI PEDIATRI DI LIBERA SCELTA AI SENSI DELL'ART. 8 DEL D.LGS. N. 502 DEL 1992 E SUCCESSIVE MODIFICAZIONI ED INTEGRAZIONI – TRIENNIO 2016-2018 [Internet]. 2022. Available from: [https://www.sisac.info/downloadFILE.do?doc_id=www.sisac.info/resources/news/news_20220428143637328&nomefile=ACN_28.04.2022_pediatri_di_libera_scelta\(2\).pdf&serv=nws](https://www.sisac.info/downloadFILE.do?doc_id=www.sisac.info/resources/news/news_20220428143637328&nomefile=ACN_28.04.2022_pediatri_di_libera_scelta(2).pdf&serv=nws) [Last accessed: 2023 May 18].
6. ACCORDO COLLETTIVO NAZIONALE PER LA DISCIPLINA DEI RAPPORTI CON I MEDICI DI MEDICINA GENERALE AI SENSI DELL'ART. 8 DEL D.LGS. N. 502 DEL 1992 E SUCCESSIVE MODIFICAZIONI ED INTEGRAZIONI – TRIENNIO 2016-2018 [Internet]. 2022. Available from: https://www.sisac.info/downloadFILE.do?doc_id=www.sisac.info/resources/news/news_20220428142843635&nomefile=ACN_28.04.2022_medicina_generale.pdf&serv=nws [Last accessed: 2023 May 18].
7. Decreto del Presidente della Repubblica 28 luglio 2000, n. 270. Regolamento di esecuzione dell'accordo collettivo nazionale per la disciplina dei rapporti con i medici di medicina generale. [Internet]. 270/2000 Jul 28, 2000. Available from: <https://www.gazzettaufficiale.it/eli/gu/2000/10/02/230/so/165/sg/pdf> [Last accessed: 2023 Jul 30].
8. Regione del Veneto. Definizione e computo dei massimali del MMG e PLS [Internet]. Available from: https://salute.regione.veneto.it/c/document_library/get_file?uuid=89897ab5-737b-4114-b13c-a40271b28e22&groupId=543512 [Last accessed: 2023 Jul 30].
9. AZIENDA USL TOSCANA NORD OVEST. OGGETTO: ACN PER LA MEDICINA GENERALE. RICOGNIZIONE INNALZAMENTO MASSIMALE INDIVIDUALE A 1800 SCELTE. [Internet]. Available from: https://www.uslnordovest.toscana.it/attachments/article/10472/20230517_1624%20decreto.pdf [Last accessed: 2023 Jul 30].
10. Programma Nazionale Esiti edizione 2022 [Internet]. Available from: https://pne.agenas.it/main/doc/Report_PNE_2022.pdf [Last accessed: 2023 May 10].
11. Guide to Prevention Quality Indicators: Hospital Admission for Ambulatory Care Sensitive Conditions. Available from: <https://www.ahrq.gov/downloads/pub/ahrqqi/pqiguide.pdf> [Last accessed: 2023 May 10].
12. TASSO DI ACCESSI IN PS NEI GIORNI FERIALE DALLE ORE 8.00 ALLE ORE 20.00 DI ADULTI CON CODICE DI DIMISSIONE BIANCO/VERDE. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_703.pdf [Last accessed: 2023 Nov 22].
13. OSPEDALIZZAZIONE PER AMPUTAZIONE DEGLI ARTI INFERIORI IN PAZIENTI DIABETICI. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_52.pdf [Last accessed: 2023 Nov 23].
14. OSPEDALIZZAZIONE PER ASMA NEGLI ADULTI. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_47.pdf [Last accessed: 2023 Nov 23].
15. OSPEDALIZZAZIONE PER BRONCOPNEUMOPATIA CRONICA OSTRUTTIVA. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_56.pdf [Last accessed: 2023 Nov 23].
16. OSPEDALIZZAZIONE PER COMPLICANZE A BREVE E LUNGO TERMINE DEL DIABETE. PROTOCOLLO OPERATIVO [Internet]. [LAST ACCESSED 2023 Nov 23]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_77.pdf [Last accessed: 2023 Nov 23].
17. OSPEDALIZZAZIONE PER DIABETE NON CONTROLLATO (SENZA COMPLICANZE). PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_49.pdf [Last accessed: 2023 Nov 23].
18. OSPEDALIZZAZIONE PER INFEZIONI DEL TRATTO URINARIO. PROTOCOLLO OPERATIVO [Internet]. Available from: <https://pne.agenas.it/assets/documentation/>

- protocolli/pro_57.pdf [Last accessed: 2023 Nov 23].
19. OSPEDALIZZAZIONE PER INFLUENZA. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_44.pdf [Last accessed: 2023 Nov 23].
 20. OSPEDALIZZAZIONE PER IPERTENSIONE ARTERIOSA. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_58.pdf [Last accessed Nov 23].
 21. OSPEDALIZZAZIONE PER SCOMPENSO CARDIACO. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_59.pdf [Last accessed Nov 23].
 22. OSPEDALIZZAZIONE PER TSO. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_702.pdf [Last accessed 2024 Jan 8].
 23. TASSO DI ACCESSI IN PS NEI GIORNI FERIALE DALLE ORE 8.00 ALLE ORE 20.00 DI PAZIENTI PEDIATRICI CON CODICE DI DIMISSIONE BIANCO/VERDE. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_704.pdf [Last accessed: 2023 Nov 23].
 24. OSPEDALIZZAZIONE PER ASMA PEDIATRICO. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_46.pdf [Last accessed: 2023 Nov 23].
 25. OSPEDALIZZAZIONE PER GASTROENTERITE PEDIATRICA. PROTOCOLLO OPERATIVO [Internet]. Available from: https://pne.agenas.it/assets/documentation/protocolli/pro_45.pdf [Last accessed: 2023 Nov 23].
 26. Ugolini C, Leucci AC, Nobilio L, Bertè G. Reorganizing territorial healthcare to avoid inappropriate ED visits: does the spread of Community Health Centres make Walk-in-Clinics redundant? *BMC Health Serv Res*. 2020 Dec;20(1):807. doi: 10.1186/s12913-020-05648-x.
 27. Lippi Bruni M, Mammi I, Ugolini C. Does the extension of primary care practice opening hours reduce the use of emergency services? *J Health Econ*. 2016 Dec;50:144–55. doi: 10.1016/j.jhealeco.2016.09.011. Epub 2016 Oct 8.
 28. Cowling TE, Harris MJ, Watt HC, Gibbons DC, Majeed A. Access to general practice and visits to accident and emergency departments in England: cross-sectional analysis of a national patient survey. *Br J Gen Pract*. 2014 Jul;64(624):e434–9. doi: 10.3399/bjgp14X680533.
 29. Cowling TE, Cecil EV, Soljak MA, Lee JT, Millett C, Majeed A, et al. Access to primary care and visits to emergency departments in England: a cross-sectional, population-based study. *PLoS One*. 2013;8(6):e66699. doi: 10.1371/journal.pone.0066699.
 30. Henricson J, Ekelund U, Hartman J, Ziegler B, Kurland L, Björk Wilhelms D. Pathways to the emergency department - a national, cross-sectional study in Sweden. *BMC Emerg Med*. 2022 Apr 7;22(1):58. doi: 10.1186/s12873-022-00619-3.
 31. Brasseur E, Gilbert A, Servotte JC, Donneau AF, D'Orio V, Ghuyssen A. Emergency department crowding: why do patients walk-in? *Acta Clin Belg*. 2021 Jun;76(3):217–23. doi: 10.1080/17843286.2019.1710040. Epub 2019 Dec 30.
 32. McIntyre A, Janzen S, Shepherd L, Kerr M, Booth R. An integrative review of adult patient-reported reasons for non-urgent use of the emergency department. *BMC Nurs*. 2023 Mar 30;22(1):85. doi: 10.1186/s12912-023-01251-7.
 33. Harris MJ, Patel B, Bowen S. Primary care access and its relationship with emergency department utilisation: an observational, cross-sectional, ecological study. *Br J Gen Pract*. 2011 Dec;61(593):e787–93. doi: 10.3399/bjgp11X613124.
 34. Safiri S, Carson-Chahhoud K, Noori M, Nejadghaderi SA, Sullman MJM, Ahmadian Heris J, et al. Burden of chronic obstructive pulmonary disease and its attributable risk factors in 204 countries and territories, 1990-2019: results from the Global Burden of Disease Study 2019. *BMJ*. 2022 Jul 27;378:e069679. doi: 10.1136/bmj-2021-069679.
 35. Steurer-Stey C, Dalla Lana K, Braun J, Ter Riet G, Puhon MA. Effects of the “Living well with COPD” intervention in primary care: a comparative study. *Eur Respir J*. 2018 Jan;51(1):1701375. doi: 10.1183/13993003.01375-2017.
 36. Jamieson AL, Harries TH, Thornton H, Crichton S, White P. Emergency Admissions for COPD in an Urban Population: The Role of Population and Primary Care Factors. *COPD*. 2015;12(6):606–12. doi: 10.3109/15412555.2015.1020150. Epub 2015 Aug 11.
 37. Pennisi F, Odelli S, Borlini S, Signorelli C, Renzi C. Impact of the Covid pandemic on timely cancer diagnosis across European healthcare settings: a scoping review. *Ann Ig*. 2024 Jan 18. doi: 10.7416/ai.2024.2596.